



CELLPHONE HAZARDS ON HEARING- AN OBSERVATIONAL STUDY IN MUMBAI POPULATION

Dr. Vaibhav Madane,¹ Dr. Nandu Kolwadkar,^{2*} Dr. Priyadershini Rangari³

¹MS (ENT), Associate Professor, Department of Otolaryngology, Government Medical College, Baramati, Pune, Maharashtra, India.

^{2*}MS (ENT), Consultant, Orange City Hospital and Research Centre, Nagpur, Maharashtra, India.

³MDS (Oral Medicine and Radiology), Assistant Professor, Department of Dentistry, Sri Shankaracharya Medical College, Bhilai, Durg, Chhattisgarh, India.

Type of Publication: Original Research Paper

Conflicts of Interest: Nil

Abstract:

Background: The mobile phone is an essential part of communication in the 21st century. However, the widespread popular use of the mobile phone device has also given rise to cautious apprehensions regarding the health hazards that might originate due to usage of the mobile phones or due to close proximity of the user to the radiofrequencies emitted from the device itself and also from the base stations.

Objectives: of the study was conducted to evaluate, compare and correlate the hearing changes in mobile phone users and non-users.

Methods: To study the impact of mobile phone usage on hearing, this study was conducted on Mumbai population, included 200 mobile phone users and non-users each, between the age group of 18-35 years.

Results: In this study, heating sensation of ears (27%) followed by blocked sensation in ears (16%), tinnitus (10%) and difficulty in hearing (4%) were the main symptoms in the study group at the time of presentation. The study and control groups were found to be comparable with respect to the sex distribution. In group 1, 5% and 3% of cases had minimal sensory-neural hearing loss in the right and left ears respectively, while in group B, 2% had sensory-neural hearing loss in the left ear only.

Conclusion: To conclude, our study did not show any significant hearing loss in mobile phone users since 0-25dB hearing loss is taken for all practical purposes as normal.

Keywords: Cell Phone, Electromagnetic Fields, Radio Frequency Radiation, Global System for Mobile Communication (GSM) Radiation, Microwaves.

Introduction

The cell phone is a useful device for communication in this modern world. If any adverse health effect is established from mobile phone use, it will be a global concern because developing countries are establishing this technology in preference to the more expensive fixed line systems. Thus, even a small impact on health could have a major public health consequence.

Mobile phones are low power radio devices that transmit and receive radio frequency radiation at

frequencies in the microwave range. The skin, inner ear, cochlear nerve and the temporal lobe surface absorb the radiofrequency energy. The extensive use of mobile phones has been accompanied by public debate on the possible adverse effects on human health. The concerns relate to the emissions of radio frequency (RF) radiation from the mobile phones and the base stations that receive and transmit the signals. There are two direct ways by which health could be affected as a result of exposure to RF radiation.¹

These are thermal (heating) effects caused mainly by holding mobile phones close to the body and also as a result of possible non-thermal effects. The well-liked belief is that adverse health effects can be induced mostly by the heating effect of Global System for Mobile Communication (GSM) radiation. The reported adverse health effects and the extensive portfolio of non-thermal effects that have been published in the scientific literature during the last few years, indicates that the kind of radiation now used in GSM phone can and does affect living organisms in various non-thermal ways. Phones emit a pulsed high-frequency electromagnetic field (PEMF)² which may penetrate the scalp and the skull. These electromagnetic fields are known to alter distinct aspects of the brain's electrical response to acoustic stimuli. The extensive exposure to microwave radiation has been found to affect a wide variety of brain functions³ such as electrical activity, electrochemistry,^{4,5} permeability of the blood brain barrier⁶ and immune system.⁷ Microwaves are known to non-thermally affect the dopamine-opiate system^{8,9} of the brain and to increase the permeability of the blood-brain barrier.¹⁰ Exposure to high-density microwaves can cause detrimental effects on the eyes, testis and other tissues and induce significant biologic changes through thermal actions.¹¹ The temporal region near the phone antenna appeared to be under the most intensive heating. Ultrahigh frequency radiation¹² induces significant changes in local temperature and in physiologic parameters of central nervous and cardiovascular systems. Besides a hypnotic effect¹³ with shortening of sleep onset latency, a REM suppressive effect with reduction of duration and percentage of REM sleep was also found. The number of complaints was higher for people using the digital (GSM) system with pulse modulated fields, than for those using the analogue Nordisk Mobile telecommunication¹⁴ (NMT) system. Radiofrequency exposure from mobile phones is concentrated to the tissue closest to the handset, which includes the auditory nerve.¹⁵ The effects on neuronal electrical activity, energy metabolism, genomic responses, neurotransmitter balance, blood-brain

barrier permeability, cognitive function, auditory function, sleep, and various brain diseases including brain tumors are of concern. Most of the reported effects are small as long as the radiation intensity remains in the non-thermal range, and none of the research reviewed gives an indication of the mechanisms involved at this range. Keeping in view the hazards of mobile phones, the present study was designed to investigate the association of use of mobile phones and hearing loss.

Materials and Methods

The prospective study was conducted in the Department of Otorhinolaryngology, Dr. D. Y. Patil Medical College, Navi Mumbai (Maharashtra), in the year March 2008 to March 2009, over a period of one year. An ethical clearance was obtained from the institutional committee prior the study. The study included 200 mobile phone users and non-users each, between the age group of 18-35 years.

The subjects selected for the study was age group between 18-35 years, >1 year of mobile phone usage, minimum usage of more than 1hour/day.

Excluded subjects those with hearing aids or implants, History of CSOM, History of head trauma or head fracture, Family history of hearing defects, People exposed to constant noise pollution/ noisy working environment.

The study was divided into two groups- Group A (cell phone users) and Group B (control group)

All candidates were subjected to a detailed history taking with special emphasis on duration, pattern, years of usage and type of mobile phone and hearing loss. Detailed enquiry was made about the onset and progression of hearing loss if any. An enquiry was made into any associated illnesses like diabetes mellitus and systemic hypertension. Any history of childhood ear discharge was asked for. The habit history was taken into consideration and occupational exposure to loud noise was also asked for and recorded. Detailed clinical examination was performed. The character of the tympanic membrane was observed in detail. The three standard tuning fork tests (Weber's, Rinne's and

Absolute Bone Conduction tests) were done. All candidates underwent a pure tone audiometry and all cases with an intact tympanic membrane underwent immittance studies. Pure tone audiograms were assessed for type and percentage of hearing loss. A detailed questionnaire for the complaints and examination findings was filled for each patient in a Performa. ENT examination was done by tuning forks—(Gardiner Brown Tuning Forks of 256Hz, 512Hz, and 1024 Hz) and Otoscopy done by a Heine Quality pocket otoscopic.

A routine Pure Tone Audiogram (0.25 to 8 kHz) was done for all patients for assessing the air conduction and bone conduction thresholds and the audiogram was plotted. For Tympanometric, the immittance test was done for all subjects using a conventional low probe tone frequency. In all cases the middle ear peak pressure, the external auditory canal volume and the peak compliance values were recorded. The primary outcome of the study was the presence or absence of hearing loss and the secondary outcomes were the presence of associated symptoms besides hearing loss amongst the subjects of the study group of mobile phone users.

Statistical analysis was done using Pearson's Chi-square test and unpaired t-test. The statistical software SPSS version 21 has been used for the analysis.

Results

The mean age of the subjects in the study and control group was 23.35 and 22.7 years respectively. The $p=0.13$ which was not statistically significant (Table 1). There were 59% males in the study group and 58% males in the control group whereas there were 41% females in the study group and 42% females in the control group. (Table 2)

Heating sensation of ears (27%) followed by blocked sensation in ears (16%), tinnitus (10%) and difficulty in hearing (4%) were the main symptoms in the study group at the time of presentation (Table: 3). None of the subjects in the control group had any symptoms at the time of presentation.

In our study group 48 (40%) subjects had 1–2 hours of exposure, 48(40%) had 2–3 hours and 24(20%) had 3–4 hours exposure to mobile phone per day (Figure 4). The $p<0.001$ which was statistically significant. (Table 3)

In our study group, 6 subjects were detected to have sensory-neural hearing loss out of which 3(6.25%) subjects used mobile phone for 2-3 hours per day and 3(12.5%) subjects used mobile phone for 3-4 hours per day for 2-3years. Mean years of exposure to mobile phones in group 1 was 2.47 years, while that of group 2 was 0.22 years with a $p<0.001$ which was statistically significant. (Table 4 and 5)

In our study group, out of 6 subjects who had hearing loss, 3 used mobile phones for 2-3 years and 3 used mobile phones for 3-4 years. Types of exposure noted in the study group were either intermittent or continuous, where continuous exposure was seen to be associated more with the minimal sensory-neural hearing loss noted in the study group, the association of which was compounded by the increasing hours and years of exposure. Exposure was of intermittent type in the control group. (Table 5 and 6)

Sensory-neural hearing loss (SNHL) was seen in 5 % (10 out of 200 subjects) of the study group and 2 % in the control group. In group A, 5% and 3% of cases had minimal SNHL loss in the right and left ears respectively, while in group B, 2% had SNHL in the left ear only. Sensory-neural hearing loss in the right ear was seen to be statistically significant with a $p<0.05$. (Table 7)

In 5% of the study group hearing loss was noted. Analysis of the data showed that the mean pure tone average in the study group was 13.7 and 14.5 dB for right and left ears respectively which was higher than the control group. (Table 8)

This indicates that although the pure tone average was within normal limits for both study and control group yet the hearing thresholds were higher for the study group and the difference in hearing thresholds between study and control group was statistically significant. It was also noted that the increase in hearing threshold

observed in the 6 subjects was in the 2kHz, 4kHz and 8kHz frequencies.

Table 1: Age distribution

Age (Years)	GROUP		P Value	Significance
	GROUP 1	GROUP 2		
	Mean ± Std. Deviation	Mean ± Std. Deviation		
	23.35 ± 2.99	22.7 ± 3.61	0.13	Not Significant

Table 2: Sex Distribution

SEX	GROUP			P Value	Significance
	GROUP 1	GROUP 2	Total		
MALE	71(59%)	70(58%)	141(59%)	>0.05	Insignificant
FEMALE	49(41%)	50(42%)	99(41%)		
Total	120(100%)	120(100%)	240(100%)		

Table 3: Symptoms on Presentation

Symptoms on presentation	Group 1	Group 2	p value
Heating sensation	27%	0%	< 0.001 (significant)
Blocked sensation	16%	0%	< 0.001 (significant)
Tinnitus	10%	0%	< 0.001 (significant)
Difficulty in hearing	4%	0%	0.024 (significant)

Table 4: Number of hours of exposure in the study group

Hours of exposure (per day)	Number of subjects	Percentage
1-2 hours	48	40%
2-3 hours	48	40%
3-4 hours	24	20%

Table 5: Years of Exposure

Years of Exposure	GROUP		P Value	Significance
	GROUP 1	GROUP 2		
	Mean ± Std. Deviation	Mean ± Std. Deviation		
	2.47 ± 0.65	0.22 ± 0.26	<0.001	Significant

Table 6: Type of Exposure

Type of Exposure	Group		Total	P Value	Significance
	Group 1	Group 2			
Intermittent	84(70%)	120(100%)	204(85)	<0.001	Significant
Continuous	36(30%)	0(0%)	36(15)		
Total	120(100%)	120(100%)	240(100)		

Table 7: Pure Tone Average Left and Right

Pure Tone Average	GROUP		P Value	Significance
	GROUP 1	GROUP 2		
	Mean ± Std. Deviation	Mean ± Std. Deviation		
Right	13.71 ± 5.61	10.86 ± 4.6	<0.001	Significant
Left	14.5 ± 5.44	11.75 ± 4.53	<0.001	Significant

Table 8: Hearing Loss

Sensorineural Hearing Loss	GROUP		Total	P Value	Significance
	Group 1	Group 2			
ABSENT	114(96)	120(100)	235(98)	0.024	Significant
PRESENT	6(4)	0(0)	5(2)		
Total	120(100)	120(100)	240(100)		

Discussion

This was an observational noninvasive study of cell phone users in the Department of Otolaryngology, Dr. D. Y. Patil Medical College, Navi Mumbai (Maharashtra).

The mean age (in years) of the subjects in the study and control group was 23.35 and 22.7 respectively. Male to female ratio in group 1 was 1.45:1 and in group 2 it was 1.4:1. The two groups were statistically insignificant with respect to the sex and age distribution (p > 0.05).

Heating sensation of ears (27%) followed by blocked sensation in ears (16%), tinnitus (10%) and difficulty in hearing (4%) were the main symptoms in the study group at the time of presentation. The control group did not present with any symptoms at the time of study. All the mentioned symptoms were found to be statistically significant.

In our study group 48(40%) subjects had 1 – 2 hours of exposure, 48(40%) had 2 – 3 hours and 24(20%) had 3 – 4 hours exposure to mobile phone per day. The p value was found to be < 0.001 which is statistically significant. Six subjects were detected to have sensory-neural hearing loss out of which 3(6.25%) subjects used mobile phone for 2-3 hours per day and 3(12.5%) subjects used mobile phone for 3-4 hours per day for 2-3years.

Mean period of exposure to mobile phones in group 1 was 2.47 years, while that of group 2 was 0.22 years with a p- value < 0.001 which was statistically significant.

Types of exposure noted in the study group were either intermittent or continuous where continuous exposure was seen to be associated more with the minimal sensory-neural hearing loss noted in the study group, the association of which was compounded by the increasing hours and years of exposure. Exposure was of intermittent type in the control group.

Sensory-neural hearing loss (SNHL) was seen in 5 % (6 out of 120 subjects) of the study group which was found to be statistically significant.

In group 1, 5% and 3% of cases had minimal sensory-neural hearing loss in the right and left ears respectively, while in group 2, 2% had sensory-neural hearing loss in the left ear only. Sensory-neural hearing loss in the right ear was seen to be statistically significant with a p value < 0.05.

Pure tone average in the study group was 13.7 and 14.5 dB for right and left ears respectively which was higher than the control group. This indicates that although the pure tone average was within normal limits for both study and control groups, yet, the hearing thresholds were higher for the study group and the difference in hearing thresholds between study and control group was statistically significant. It was also noted that the increase in hearing threshold observed in the 6 subjects was in the 2 kHz, 4 kHz and 8 kHz frequencies.

In this study, heating sensation of ears (27%) followed by blocked sensation in ears (16%), tinnitus (10%) and difficulty in hearing (4%) were the main symptoms in the study group at the time of presentation. None of the subjects in the control group had any symptoms at the time of presentation. In contrast, in a study conducted by Thahir I-Khlaiwi et al^[32], the overall mean percentage for presenting complaints in all groups were headache (21.6%), sleep disturbance (4%), tension (3.9%), fatigue (3%) and dizziness (2.4%). Sultan A Meo SA et al^[34] in a study showed about 34.59% of problems were related with impaired hearing, ear ache and/or warmth on the ear, and 5.04% of complaints with the decreased and/or blurred vision.

In our study, 3 of those who have been using mobile phone since 2 years had SNHL (sensory-neural hearing loss) and 3 of those who have been using it since 3 years had SNHL. The minimal hearing loss noted in the study group depended not only on years of exposure, but also on the hours of exposure per day and type of exposure and it was noted that with the increase in hours of exposure per day and years of exposure the incidence of hearing loss also increased. Naresh K. Panda et al^[34] found no significant difference was found for high frequency hearing loss,

DPOAE (distortion product oto-acoustic emissions), ABR (auditory brain stem response) and MLR in the users. High frequency hearing loss was seen in subjects using the phone for more than 4 years (P= 0.040).

In our study, we also observed that none of the subject in the study group suffered from significant hearing loss (0-25dB hearing loss is taken for all practical purposes as normal). This correlates with the findings of Harry C Davidson H C et al^[34]. High or long-term users reported no worse hearing, tinnitus, or balance than low or short-term users. However there were no harmful effects of mobile phone usage on their audio vestibular systems within the range of exposure of the study, in so far as can be detected by the self-report method employed.

Similarly, in a study done by Uwe Sievert et al^[35] to investigate the biological effect of the high frequency radiation produced by the Global System for Mobile Communications (GSM) mobile phone on the inner ear with its sensors of the vestibular and auditory systems showed that the electromagnetic fields generated in using the mobile phone do not have an effect on the inner ear and auditory system to the colliculus inferior in the brainstem and on the vestibular receptors in the inner ear and the vestibular system.

Although none in our study group had significant hearing loss, we observed an increase in hearing threshold between 5-15dB in the study group especially in those exposed to mobile phone usage for more than 2 hours/day, more so if the pattern of usage is continuous. This relates to the study done by García Callejo FJ et al^[37] he found audiometric curve was similar in cases and controls at the beginning of the study. After this follow-up, cases showed an increase on hearing threshold between 1 and 5 dB HL more than controls in speech tones (p<0.001).

Our findings were also in agreement with the study done by M Faruk Oktay et al^[37] Brainstem evoked response audiometric (BERA) results showed no differences among the groups (p > 0.05). In PTA measurements, no differences were observed between moderate mobile phone users (10-20 min. per day) and control subjects.

However, detection thresholds in those who talked approximately 2 h per day were found to be higher than those in either moderate users or control subjects. However, detection thresholds in those who talked approximately 2 h per day were found to be higher than those in either moderate users or control subjects. Differences at 4000 Hz for both bone and air conduction for right ears, and 500 Hz, and 4000 Hz bone and air conduction for left ears were significant for mean hearing threshold. This study shows that a higher degree of hearing loss is associated with long-term exposure to electromagnetic (EM) field generated by cellular phones.

A case of sensory-neural hearing loss due to Global System for Mobile Communications mobile was reported in Saudi medical journal in 2007.^[39]

While, those in our study with mobile phone usage of less than 2 hour/day showed normal audiometric curves which can be related to study done by Uwe Sievert et al^[40] in 12 healthy test persons, with normal hearing. Auditory brain stem reflexes recordings were performed before, during, and after exposure to electromagnetic emissions by standardized mobile phone devices. Two modes of electromagnetic emissions fields were administered: pulsed and continuous. For acoustic stimulation simultaneous to field exposure, special "plug-in" earphones had to be used. There are no adverse effects of mobile phone emissions on the ear function, at least on a short-term range.

Conclusion

To conclude, our study did not show any significant hearing loss in mobile phone users since 0-25dB hearing loss is taken for all practical purposes as normal. But still we observed a minimal hearing loss of 5-15dB in those exposed to mobile phone usage of more than 2 hours per day, the pattern of use being continuous. Also noted was that the threshold was raised for the high frequencies which are not calculated under routine pure tone average. Hence we recommend a long term follow up study in long term mobile users before we can come to a definitive conclusion.

References

1. Thamir I, Sultan A Meo. Saudi Medical Journal 2004; Vol. 25 (6): 732-36.
2. Eulitz C, Ullsperger P, Freude G, Elbert T. Mobile phones modulate response patterns of human brain activity. Neuro Report 1998; 9: 3229-32.
3. Reiser H, Dimpfel W, Schober F. The influence of electromagnetic fields on human brain activity. Eur J Med Res 1995; 1: 27-32.
4. Dutta SK, Subramoniam A, Ghosh B, Parshad R. Microwave radiation-induced calcium ion efflux from human neuroblastoma cells in culture. Bioelectromagnetics 1984; 5: 71-78.
5. Frey AH. Electromagnetic field interactions with biological systems. FASEB J 1993; 7: 272-81
6. Salford LG, Brun A, Sturesson K, Eberhardt JL, Persson BR. Permeability of the blood-brain barrier induced by 915 MHz electromagnetic radiation, continuous wave and modulated at 8, 16, 50, and 200 Hz. Microsc Res Tech 1994; 27: 535-42.
7. Youbicier-Simo BJ, Boudard F, Cabaner C, Bastide M. Biological effects of continuous exposure of embryos and young chickens to electromagnetic fields emitted by video display units. Bioelectromagnetics 1997; 18: 514-23.
8. Becker RO, Marino AA. Electromagnetism and Life. Albany (NY): Suny Press; 1982.
9. Frhlich H. Advances in Electronics and Electron. Physics 1980; 53: 85-152.
10. Bawin SM, Kaczmarek LK, Adey WR. Effects of modulated VHF fields on the central nervous system. Ann N Y Acad Sci 1975; 247: 74-81.
11. Nakamura H, Matsuzaki I, Hatta K, Nobukuni Y, Kambayashi Y, Ogino K. Non-thermal effects of mobile-phone frequency microwaves on utero-placental functions in pregnant rats. Reprod. Toxicol 2003; 17: 321-26.
12. Khudnitskii SS, Moshkarev EA, Fomenko TV. On the evaluation of the influence of cellular phones on their users. Med. Tr Prom Ekol 1999; 9: 20-24.

13. Mann, K, Roschke, J, Effects of pulsed high-frequency electromagnetic fields on human sleep. *Neuro-psychobiology* 1996; 33: 41-47.
14. Sandstrom M, Wilen J, Oftedal G, Hansson MK. Mobile phone use and subjective symptoms. Comparison of symptoms experienced by users of analogue and digital mobile phones. *Occup. Med* 2001; 51: 25-35.
15. Stefan Lonn, Anders Ahlbom, Per Hall, Maria Feychting. Mobile phone use and the risk of acoustic neuroma. *Epidemiology*: November 2004; 15(6).p.653-59.
16. J Ballyntine. Anatomy of the ear. In: Ballyntine J, Groves J, editors. *Scott-Brown's diseases of the ear, nose and throat*. 4th ed. London: Butterworths; 1979. p. 61-107.
17. Bitner-Glindzicz M. Hereditary deafness and phenotyping in humans. *Br Med Bull* 2002; 63:73-94.
18. Occupational Safety and Health. *Scientific American Medicine*. CTM. 1991 Chapter VII.p.18.
19. Burns W. Noise and Man. In: John Murray, editor. 2nd ed. London: 1973. p. 190.
20. J Ballantyne, J Groves. *Scott-Brown's Diseases of the ear, nose and throat*. 4th ed. London: Butterworths; 1979. Vol 2. p. 568.
21. Ward WD. Noise Induced Hearing Damage. In: Paparella MM, Shumrick DA, editors. *Otolaryngology*. 2nd ed. 1980. Vol. 2. p. 1789.
22. Snow JB. Management and therapy of trauma to the external ear and auditory and vestibular systems. In: Alberti PW, Ruben RJ, editors. *Otologic Medicine and Surgery*. New York: Churchill Livingstone; 1988. Vol 2. p. 1568.
23. Edward Arnold. In: Harold Ludman, editor. *Mawson's Diseases of the Ear*. 5th ed. London: 1988. p. 615-16.
24. Schweitzer VG. Ototoxicity of chemotherapeutic agents. *Otolaryngology Clinics of North America* 1993; 26(5): 759-89.
25. Martindale The extra pharmacopoeia. In: James E. F. Reynolds, editor. 29th ed. The Pharmaceutical Press: London. p. 508-18.
26. Edward Arnold. In: Harold Ludman, editor. *Mawson's Diseases of the Ear*. 5th ed. London: 1988. p. 617.
27. Brummett RE. Ototoxic liability of erythromycin and analogues. *Otolaryngology Clinics of North America* 1993; 26(5): 811-19.
28. Nadol JB. Hearing loss (Review). *The New England Journal of Medicine* 1993; 329(15): 1092-102.
29. Ballenger JJB. Diseases of the Nose, Throat, Ear, Head and Neck. In: Lea, Febiger, editors. 14th ed. London: 1991. p. 1209.
30. Edward Arnold. In: Harold Ludman, editor. *Mawson's Diseases of the Ear*. 5th ed. London: 1988. p. 397-98.
31. Luigi Parmeggiani, editor. *Encyclopaedia of Occupational Health and Safety*. 3rd (revised) ed. International Labour Office: Geneva: 1983. P. 596.
32. Thamir I-Khlaiwi, Sultan A Meo. *Saudi Medical Journal* 2004; Vol. 25 (6): 732-36.
33. Sultan A. Meo, Abdul M Al-drees. Mobile phone related-hazards and subjective hearing and vision symptoms in the Saudi population. *International journal of occupational medicine and environmental health (Int J Occup Med Environ Health)* 2005; vol. 18 (issue 1). p. 53-7.
34. Naresh K. Panda, Rishab Jain, Jaimanti Bakshi. Audiological Disturbances in Long-Term Mobile Phone Users. *Otolaryngology-Head and Neck surgery*. August 2007; Vol 137 (Issue 2) Supplement. p.131-32.
35. Harry C Davidson, Mark ELutman. Survey of mobile phone use and their chronic effects on the hearing of a student population. *International journal of audiology (Int J Audiol)* 2007 Mar; vol. 46 (issue 3). P. 113-18.
36. Uwe Sievert, Siegfried Eggert, S Goltz, Hans Wilhem. Effects of electromagnetic fields emitted by cellular phone on auditory and vestibular labyrinth. *Laryngo- rhino- otologie (Laryngorhinootologie)* 2007 Apr; vol. 86 (issue 4). p. 264-70.
37. García Callejo FJ, Peña Santamaría. Hearing level and intensive use of mobile phones.

- Acta otorrino laringológica española (Acta Otorinolaringol Esp). 2005 May; vol. 56 (issue 5) .p. 187-91.
- 38.** M FarukOktay, Suleyman Dasdag. Electromagnetic biology and medicine (Electromagn. Biol. Med). 2006; vol. 25 (issue 1). P. 13-21.
- 39.** Saudi medical journal (Saudi Med J), 2007 Aug; vol. 28 (issue 8). P. 1283-6.
- 40.** Uwe Sievert, Siegfried Eggert, Hans Wilhem. Can mobile phone emissions affect auditory functions of cochlea or brain stem? Otolaryngology—Head and Neck Surgery, Otolaryngol Head Neck Surg. 2005, 132: 3: 451-5.
- 41.** Parazzini M, Brazzale AR, Paglialonga A, et al. Effects of GSM cellular phones on human hearing: the European project "GUARD". Radiat Res, 2007 Nov; 168: 5: 608-13.
- 42.** Uloziene I, Uloza V, Gradauskiene E; Saferis V. Assessment of potential effects of the electromagnetic fields of mobile phones on hearing. BMC Public Health, 2005, 5: 1: 39.
- 43.** Kerekhanjanarong V, Supiyaphun P, Naratricoorn J, Laung pitackchumpon P. The effect of mobile phone to audiologic system. J Med Assoc Thai. 2005; 88 Suppl 4:S231-4.
- 44.** Balbani AP, Montovani JC. Mobile phones: influence on auditory and vestibular systems. Braz J Otorhinolaryngol. 2008; 74:1:125-31.
- 45.** Ramya C S, Karthiyanee K, Vinutha S. Effect of mobile phone usage on hearing threshold: A pilot study. Indian J Otol 2011; 17:159-61.